

**CLASSIFICATION OF DRAINAGE BASINS FOR ENVIRONMENTAL PURPOSES IN  
GIS PLATFORM USING SOFT COMPUTING APPROACH**

**GOURNELOS, TH.<sup>1</sup>, EVELPIDOU, N.<sup>2</sup>, VASSILOPOULOS, A.<sup>2</sup>**

**ABSTRACT**

The drainage basin development reflects the dominant erosional processes and its internal rock structure. Many variables control the drainage basin evolution: lithology, topography, geology and climatic conditions.

The purpose of this paper is to propose different classification mechanisms of the drainage basins, based on the above controlling variables. These classification schemes can be divided in two categories: the Boolean model and the Fuzzy model, depending of the boundary vagueness of the initial data.

Such classification of drainage basins can become very useful for developing environmental protection strategies. Some examples are given from the Greek drainage system.

**KEYWORDS:** Erosion Risk, Classification methods, GIS, Geomorphology, Thera

**INTRODUCTION**

The geomorphological processes are being controlled by many variables. Firstly weathering of the rocks occurring by mechanical or chemical factors, which disintegrate or decomposes chemically the rocks. Next erosional processes are taking place. This depends on various factors such as the lithology, the slope gradient, the vegetation cover, the climatic conditions, the structure of the rocks and the erosivity of the erosion agent. The denudation of the existing landforms is mainly achieved by river or channel systems. Thus the drainage basin is considered a fundamental unit for studying such geomorphological process.

---


1:Associate Professor, University of Athens, Geology Department, Physical Geography & Climatology Sector, Papepistimiopolis, Zografou, 157-84.

2:Dr. Geologist, University of Athens, Geology Department, Physical Geography & Climatology Sector, Papepistimiopolis, Zografou, 157-84.

## METHODOLOGY

The classification of drainage basins or sub-basins evaluating erosion risk of soil or rocks involves a multi-step procedure. Firstly input data from topographical, geological, and land use maps, aerial photos interpretation and field observations are introduced in a GIS system. The scale of the input data depends on the needed detail of the derived data. The necessary input variables from the above data, sources are contour lines, geomorphological characteristics such as drainage system, geological formations and land use / land cover. Processing the above data secondary variables can be produced such as morphological slopes, drainage density and vulnerability of the rocks.

Following, a first set of logical rules is applied to the main drainage basins achieving a first level classification. These rules are usually of a fuzzy nature (Zadeh, L.A., 1965, Zadeh, L.A., 1987) because of the relative imprecision of the involved variables such as rocks erodibility, slope gradient, drainage density, etc.




After the first classification an output map is created in GIS, showing the spatial distribution of basins with different degrees of erosion risk. The drainage basins which present high or very high degrees of erosion risk are further decomposed using a new set of logical rules, and thus a second level of classification of drainage sub-basins is achieved. Finally the same procedure is

repeated to arrive in the third level of classification of drainage sub-set of sub-basins (Fig. 1).


### CASE STUDY: THE ISLAND OF THERA

The island of Thera is chosen for applying the above methodology (Fig. 2), for several reasons. Firstly the geological evolution of this island and the nature of



surface sediments render Thera very susceptible to geomorphological processes. Secondly an extensive work has been done the previous years (Gournelos et al, 1995, Gournelos et al, 1999) by the authors, concerning the geomorphology of the island.

This island consists of a substratum of recrystallized limestones and metamorphosed rocks and a sequence of volcanic formations (Papastamatiou, 1958, Tataris, 1964, Fytikas et al, 1984, Friedrich and Velitzelos 1986, Skarpelis and Liati, 1987, Astaras et al, 2001). This volcanic activity started about 1,6 m.y. and its last phase was dominated by the Minoan explosion (Galanopoulos, 1958, Seward et al, 1980, Pichler and Kussnaul, 1980, Druit et al, 1989). Thus the Minoan eruption is responsible for the present landforms of the islands and the surface's rocks (Upper-Pumice series).




*Fig. 2: The location of Thera island and a view of its morphology.*

The input data that were used in this case study, were the contours with interval of 50m, deriving from

the digitization of topographical maps scale 1:50.000 (HAGS, 1970, Topographical map, Thera Sheet, scale 1:50.00), the geological formations and tectonical structure from geological maps scale 1:50.000 (IGME, 1980, Geological map, Thera

Sheet, scale 1:50.000), the land use from different kind of land use maps (Ministry of Agriculture, 1986, Astaras, Th., et al, 1998) and from interpretation of aerial photos, scale 1:30.000 (HAGS, 1988, scale 1:30.000). Figure 3 shows some of the input data from which secondary information such as morphological slopes, drainage density and frequency have been derived.

The estimation of the rock's vulnerability was based on Selby's (1987) proposal, which considers the rock mass strength classification and rating, to



express the resistance to erosion. Using this kind of classification limestones are more resistant to erosion than schists. The first attempt to assign erodibility values on different rock types was made by Jensen and Painter, (1974).

The classifications procedure for the Thera island is shown in figure 4 and is as follows:

- From the input data (geological, lithological formations, tectonical structures, topographical and morphological characteristics) the rocks erodibility was estimated. The erodibility of the rocks was the only variable, used in the first




Fig. 4: Classification flow diagram that used in Thera case study.




Fig. 5: First classification level, showing that the whole island except the 'caldera' area is considered as high erosion risk.

classification. The main geomorphological units are the caldera with a series of lavas and pumice deposits and the rest of the island, which was dominated by the Minoan pumice series. Figure 5 presents the first classification of Thera island. As shown the first classification concluded that the whole island except the caldera part is characterized of high erosion risk. The area outside of the drainage basins has not been calculated.

- The rest of the 'caldera' part, island, which has been indicated by the first classification as of

high erosion risk, is further classified applying logical rules such as:

*"If the erodibility of the rocks is High and slope gradient is High then Erosion risk is Very High".*

The results of the second classification are shown in the figure 6. As shown in the legend of this figure, 42 drainage basins are of high erosion risk, meaning that the 53,16% of the studied drainage basins are of high erosion risk, while only the 5,06% and 16,45% are of low and very low erosion risk correspondingly.

- Following, the drainage basins of high and very high erosion risk indicated by the second classification have been chosen for the third level classification. The third level classification is achieved using the same logical rules as those in the second level classification. The results of the third classification are shown in Figure 7.

## **CONCLUSIONS**

In this paper different schemes are proposed for classification of drainage basins or subsets of drainage basins. The most useful model is the fuzzy one, because relative imprecision of distribution values characterizes the involved variables. We applied this methodology in the case study of the Thera island and a series of output maps is produced. These maps show spatial distribution of the erosion risk degree of the areas of Thera in different scales and can be used in local planning for environmental management.

## **REFERENCE**

- [1]Astaras, T., Lambrinos, N. and Soulakelis, N., 2001, Multitempotal monitoring at Santorini volcano by use of short and thermal infrared LANDSAT-5 TM Images, Proceedings International symposium on engineering geology and the environment, Vol. 3, p. 495-503.
- [2]Astaras, Th., Lambrinos, N. and Manakos, Y., 1998, Integration of Remote Sensing data in GIS for assessment of land complexes of Santorini island, Proceedings of the second workshop, Greece, 259-269.
- [3]Druitt, T.M., Mellors, R.A., Pyle, D.M., and Sparks, R.S.J., 1989, Explosive volcanism on Santorini, Greece, Geol. Mag. 126, 95-126
- [4]Friedrich, W. and Velitzelos, E., 1986, Bemerkungen zur spatquartaren Flora von' Santorini/Griechenland, Courier Forschungs, Inst. Senckenberg 86, 387-395.
- [5]Fytikas, M., Innocenti, F., Manetti, P., Mazzuoli, R., Peccerillo, A. and Villari, L., 1984, Tertiary to Quaternary evolution of volcanism in the Aegan region, in J.E. Dixon and A.H.F. Robertson (eds.), The Geological Evolution of the Eastern Mediterranean (Geol. Soc. London Spec. Publ. 17), 687-699
- [6]Galanopoulos, A. G., 1958, Zur Bestimmung des Alters de Santorini - Kaldera. Annales geol. Pays hell, 9, 185-188
- [7]Gournelos, Th., Vaiopoulos, D., Vassilopoulos, A. and Evelpidou, N., 1995, Geomorphological study of Thera island, Palaio-anaglyf synthesis, Proceedings of the 4th Hellenic Geographical Congress.
- [8]Gournelos, Th., Vassilopoulos, A. and Evelpidou, N., 1999, Erosion risk study in GIS environment using fuzzy theory, Proceedings of the 1st Hellenic congress:

Geographical Information Systems.Capabilities and Invitations, CD-Rom.

- [9]HAGS (Hellenic Army Geographical Service), 1975, Topographic Map, Thera sheet, Scale 1:50.000, Athens.
- [10]HAGS (Hellenic Army Geographical Service), 1988, Aerial photos, Thera island, Scale 1:30.000, Athens.
- [11]IGME (Institute of Geology and Mineral Exploration), 1980, Geological Map, Thera sheet, Scale 1:50.000, Athens.
- [12]Jensen, J.M. and Painter, R.B., 1974, Predicting sediment yield from climate and topography, J. Hydrol., v.21, p.p. 371-380.
- [13]Ministry of Agriculture, 1986, Soil Map of Greece, Thera sheet, Scale 1:50.000.
- [14]Papastamatiou, I.N., 1958, The age of crystalline limestones of Thera island, Jour Geol. Soc. Greece III-1, p.p. 104-113.
- [15]Pichler, H. and Kussmaul, S., 1980, Comments on the geological map of the Santorini Islands, in C. Doulas (ed.), Thera and the Aegan World II, 413-427
- [16]Selby, M.J., 1987, Rock slopes. In Anderson, M.G., and Richards, K.S., (eds), Slope Stability, Wiley, Chichester, 475-504.
- [17]Seward, D., Wagner, G. and Pichler, H., 1980, Fission-track ages of Santorini volcanics (Greece), in C. Doulas (ed.), Thera and the Aegan World II, 101-1108
- [18]Skarpelis, N. and Liati, A., 1987, Granite intrusion, skarn formation and mineralization in the metamorphic basement of Thera, Cyclades, Greece, 5th meeting European Geological Societies, Dubrovnic.
- [19]Tataris, A.A., 1964, The presence of Eocene in the semimetamorphic basement on Thera's island, Jour. Geol. Soc. of Greece VI-1, p.p. 232-236.
- [20]Zadeh, L.A.: 1965, Fuzzy sets, Information and Control, 8, p.p. 338-353.
- [21]Zadeh, L.A.: 1987, The concept of linguistic variable and its application to approximate reasoning, R.R. Yager, S. Ovchinnikov, R.M. Tong, H.T. Nguyen (eds), Fuzzy Sets and Applications, Wiley, New York, p.p.293-329.




Fig.6: Erosion risk map of 2<sup>nd</sup> level drainage basins classification.




Fig.7: Erosion risk map of 3<sup>rd</sup> level drainage sub-basins classification.