

The 25th October, 2018 Zakynthos Earthquake

P. Papadimitriou¹, V. Kapetanidis¹, A. Karakonstantis¹, I. Spingos¹, K. Pavlou¹, G. Kaviris¹, I. Kassaras¹ and N. Voulgaris¹

(1) Section of Geophysics–Geothermics, Department of Geology and Geoenvironment, National and Kapodistrian University of Athens, 15784 Panepistimiopolis, Zografou, Athens, Greece, ppapadim@geol.uoa.gr

The geodynamic regime in western Greece manifested by intense seismic activity results from the convergence between the African and Eurasian lithospheric plates (e.g. Armijo *et al.*, 1992), while the westward motion of the Turkish plate imposes an important contribution (e.g., Armijo *et al.*, 1996). The region hosts both continental collision to the north and oceanic slab subduction to the south; these features are connected by the right-lateral Cephalonia Transform Fault (CTF). Active tectonics involves all types of faulting. Reverse and strike-slip faulting characterize the interplate type of deformation (Papadimitriou *et al.*, 2006; 2012; Kassaras *et al.*, 2016), while normal and strike-slip faulting prevails in intraplate regions (Kassaras *et al.*, 2014a, 2014b), forming pull-apart basins (King *et al.*, 1993) and the rapidly spreading Corinth Rift (Kapetanidis & Papadimitriou, 2011; Kapetanidis *et al.*, 2015). GPS measurements demonstrate rapid deformation to the south (~ 35 mm-yr⁻¹) and a slower one of the order 5–8 mm-yr⁻¹ in NW Greece (Chousianitis *et al.*, 2015). Deformation across the area of Western Greece is mainly accommodated within the accretionary sediments comprising the external Paxi, Gavrovo and Ionian geotectonic units, escarped from the African margin and colliding against the un-deformed Eurasian hinterland (e.g. Royden and Papanikolaou, 2011).

Zakynthos Island is located at the front of the present-day Hellenic Trench, which is formed along the convergent zone of the plate boundaries between the subducting oceanic lithosphere of Tethys and the overriding Eurasian plate. The high level of seismic activity in the area is the result of the intense crustal deformation in the Central Ionian Sea, which leads to the occurrence of moderate to strong events. The most recent events include the November 18th, 1997 (Mw=6.5) earthquake, SW of Zakynthos Island, the December 2nd, 2002 (Mw= 5.5) earthquake, a sequence of earthquakes on October 2005 (Mw=5.6) and April 2006 (Mw=5.5–5.7) south of Zakynthos, the June 8th, 2008 (Mw= 6.4) Andravida earthquake, the January 26th, 2014 (Mw= 6.1) Paliki earthquake and the November 17th, 2016 (Mw= 6.4) Lefkas earthquake.

A strong earthquake of moment magnitude Mw=6.8 occurred on 25 October 2018, ~ 50 km to the SW of Zakynthos Island, causing some damage in the municipality of Zakynthos and on the islands of Strofades. This event was one of the largest that have occurred in the vicinity of Zakynthos Island since 1983. It was strongly felt in Western Greece but also in other parts of the Greek mainland. More than 4,000 aftershocks were located in the framework of this study. The aftershock zone covers a region about 60-70 km long in a N110°E direction and ~ 80 km towards N20°E.

In this work we attempt to highlight the nature and dynamics of the earthquake sequence, the driving forces that acted during its evolution and potential consequences of its occurrence on the regional hazard. To this aim, a comprehensive dataset of recordings from the Hellenic Unified Seismological Network (HUSN) and the Hellenic Strong Motion Network (HSMN; Theodulidis *et al.* 2004) was compiled and the following tasks were carried out: (a) determination of precise hypocentral locations employing a custom velocity model, followed by double-difference relocation, (b) spatiotemporal analysis of the sequence, (c) regional moment tensor inversion for the determination of the source parameters of the mainshock and the larger events, (d) inversion of focal mechanisms to investigate the local stress-field distribution and (e) computation of Coulomb stress transfer to identify regions and receiver faults that were loaded with additional stress.

After the manual determination of P- and S-wave phase arrivals, a dataset of 900 best located earthquakes, between 26 October and 30 November 2018, was compiled and a local velocity model was obtained using a travel-time residual and location errors minimization method (e.g. Kissling *et al.*, 1994). Locations were recalculated with the new model and relocated with the use of a double-difference algorithm. The spatial distribution of epicenters revealed several distinct clusters, some of which are in general agreement with the geometry of known mapped faults and compatible with the strike of Quaternary faults.

Regional moment tensor inversion revealed a focal mechanism with a strike of N16°E, a dip of 25° and a rake of 166°. Most of the determined focal mechanisms of the sequence followed this trend. Nevertheless, the largest aftershock displayed strike-slip characteristics. In addition, focal mechanisms, including events of past activity (Kassaras *et al.*, 2016), were inverted to determine the regional stress field (Hardebeck and Michael, 2006). A complex stress field in the broader area of Central Ionian Sea, related to both thrust and strike-slip faulting was revealed.

Acknowledgements

We acknowledge support of this study by the project “HELPOS – Hellenic Plate Observing System” (MIS 5002697) which is implemented under the Action “Reinforcement of the Research and Innovation Infrastructure”, funded by the Operational Programme “Competitiveness, Entrepreneurship and Innovation” (NSRF 2014-2020) and co-financed by Greece and the European Union (European Regional Development Fund).

References

- Armijo, R., Lyon-Caen, H., Papanastassiou, D., 1992. East-west extension and Holocene normal-fault scarps in the Hellenic arc. *Geology* 20, 491–494.
- Armijo, R., Meyer, B., King, G.C.P., Rigo, A., Papanastassiou, D., 1996. Quaternary evolution of the Corinth Rift and its implications for the Late Cenozoic evolution of the Aegean. *Geophys. J. Int.* 126, 11–53.
- Chousianitis, K., Ganas, A., Evangelidis, C.P., 2015. Strain and rotation rate patterns of mainland Greece from continuous GPS data and comparison between seismic and geodetic moment release. *J. Geophys. Res. Solid Earth* 120, 3909–3931.
- Hardebeck, J.L. and Michael, A.J., 2006. Damped regional-scale stress inversions: Methodology and examples for southern California and the Coalinga aftershock sequence. *J. Geophys. Res. Solid Earth*, 111, B11310.
- Kapetanidis, V., Papadimitriou, P., 2011. Estimation of arrival-times in intense seismic sequences using a Master-Events methodology based on waveform similarity. *Geophys. J. Int.* 187, 889–917.
- Kapetanidis, V., Deschamps, A., Papadimitriou, P., Matrullo, E., Karakonstantis, A., Bozionelos, G., Kaviris, G., Serpetsidaki, A., Lyon-Caen, H., Voulgaris, N., Bernard, P., Sokos, E., Makropoulos, K., 2015. The 2013 earthquake swarm in Helike, Greece: seismic activity at the root of old normal faults. *Geophys. J. Int.* 202, 2044–2073.
- Kassaras, I., Kapetanidis, V., Karakonstantis, A., Kaviris, G., Papadimitriou, P., Voulgaris, N., Makropoulos, K., Popandopoulos, G., Moshou, A., 2014a. The April-June 2007 Trichonis Lake earthquake swarm (W. Greece): New implications toward the causative fault zone. *J. Geodyn.* 73, 60–80.
- Kassaras, I., Kapetanidis, V., Karakonstantis, A., Kouskouna, V., Ganas, A., Chouliaras, G., Drakatos, G., Moshou, A., Mitropoulou, V., Argyrakis, P., Lekkas, E., Makropoulos, K., 2014b. Constraints on the dynamics and spatio-temporal evolution of the 2011 Oichalia seismic swarm (SW Peloponnesus, Greece). *Tectonophysics* 614, 100–127.
- Kassaras, I., Kapetanidis, V. and Karakonstantis, A., 2016. On the spatial distribution of seismicity and the 3D tectonic stress field in western Greece. *Phys. Chem. Earth, Parts A/B/C*, 95, 50–72.
- King, G., Sturdy, D., Whitney, J., 1993. The landscape geometry and active tectonics of northwest Greece. *Geol. Soc. Am. Bull.* 105, 137–161.
- Kissling, E. et al., 1994. Initial reference models in local earthquake tomography. *J. Geophys. Res.* 99: 19635-19646.
- Papadimitriou, P., G. Kaviris and K. Makropoulos, 2006. The Mw=6.3 2003 Lefkada Earthquake (Greece) and induced transfer changes. *Tectonophysics*, 423, 73-82.
- Papadimitriou, P., Chousianitis, K., Agalos, A., Moshou, A., Lagios, E. and Makropoulos, K., 2012. The spatially extended 2006 April Zakynthos (Ionian Islands, Greece) seismic sequence and evidence for stress transfer. *Geophysical Journal International*, 190 (2), 1025-1040.
- Royden, L.H., Papanikolaou, D.J., 2011. Slab segmentation and late Cenozoic disruption of the Hellenic arc. *Geochemistry, Geophys. Geosystems* 12(3).
- Theodulidis N. et al., 2004. HEAD1.0 : A unified accelerogram database”, *Seism. Research. Lett*, 75, 41-50.