

# The observability of plasmoid-powered $\gamma$ -ray flares w the *Fermi* Large Area Telescope



Maria Petropoulou


Department of Physics  
National & Kapodistrian University of Athens

In collaboration with:


Manuel Meyer (Erlangen Center for Astroparticle Physics)  
Ian Christie (Northrop Grumman)

on behalf of the *Fermi*-LAT Collaboration

# Blazars





- Most extragalactic  $\gamma$ -ray sources are active galactic nuclei (AGN)
- Blazars are AGN with jets closely aligned to the line of sight




# Motivation

- Blazar  $\gamma$ -ray variability occurs on different timescales (months/weeks to hours/minutes)
- Short-duration flares usually emerge on top of slower varying emission
- Origin of short-duration flares is still a mystery!


Shukla & Mannheim (2020)



Meyer, Scargle, Blandford (2019)



# Magnetic reconnection




Credit: H. Hakobyan

When?

Magnetic field lines of opposite polarity are brought together by bulk plasma motions

What?

Magnetic energy is transformed to heat, bulk plasma kinetic energy and kinetic energy of relativistic particles




Credit: H. Hakobyan

Current sheet fragments into quasi-spherical plasma structures known as *plasmoids* that

- contain relativistic particles in rough energy equipartition with magnetic fields
- can move along the current sheet with speeds approaching the speed of light
- have a finite lifetime

Each plasmoid produces a flare with characteristic duration and flux !

# Model





# Model parameters

	Model			
	A	B	C	D
Bulk Lorentz factor, $\Gamma_j$	12	24	24	24
Observer's angle, $\theta_{\text{obs}}$ (deg)	0	0.2	0	0
Angle between layer and jet axis (in jet frame), $\theta'$ (deg)	0	0	30	0
Dissipation distance, $z_{\text{diss}}$ (pc)	0.2	0.4	0.4	0.4
Bolometric luminosity of external radiation, $L_{\text{ext}}$ ( $10^{45}$ erg $\text{s}^{-1}$ )	4	4	4	10
Luminosity of two-sided jet, $L_j$ ( $10^{47}$ erg $\text{s}^{-1}$ )	1	5	5	5
Target blazar	3C 273	3C 273	3C 273	3C 279


Fixed parameters (adopted from Christie et al. 2019)

Half-length of current sheet	Plasma magnetization	Magnetic field strength at $z_{\text{diss}}$	Average co-moving energy density of relativistic pairs	Average minimum Lorentz factor of pairs	Average maximum Lorentz factor of pairs	Power-law slope of pair distribution at injection
$5 \cdot 10^{16}$ cm	10	5 G	$2.2 \text{ erg cm}^{-3}$	94	5000	2.1

# Theoretical light curves

Luminous ultra-rapid flares  
from fast plasmoids

Long-duration flare from slow  
"monster" plasmoid



Variability sensitive to orientation of  
observer and current sheet



# Goals

- Evaluate which features of theoretical light curves (e.g. ultra-rapid variability) could be detected with Fermi-LAT
- Check if a quantitative comparison of the model to the data is feasible



# Methods

- Simulate artificial LAT light curves for **3C 273** and **3C 279**
- Simulation and analysis done with `fermipy`:
  - Perform standard analysis for each light curve bin
  - Replace central source of best-fit model of region of interest (ROI) with reconnection model prediction, multiply with EBL and BLR optical depth, add quiescent source flux
  - Re-run simulation of modified ROI and LAT analysis





# Artificial LAT light curves

Ultra-rapid flares are washed out due to binning

### 3 hr binned light curves

Flares in artificial binned light curves are the superposition of many plasmoid flares


Average flux level of bright 3C 273 flares is reproduced



Difficult to match the flux of the brightest flare of 3C 279

# Artificial LAT light curves: fast variability

## 5-min binned light curves







Ultra-rapid flares are detectable, if these occur within GTIs !

# Artificial LAT light curves: periodograms

$$\text{PSD}(v) \sim v^{-k}$$

Model	$k_{\text{art}}$	$k_{\text{obs}}$
A	$< 1.47$	$1.18^{+1.82}_{-0.97}$
B	$< 0.70$	$1.18^{+1.82}_{-0.97}$
C	$< 2.56$	$1.18^{+1.82}_{-0.97}$
D	$< 0.83$	$1.67^{+1.33}_{-1.34}$




PSD slopes of  
real & artificial  
light curves are  
compatible  
within errors



Hard PSDs (slope  $\sim 0$ ) in  
probed frequency range

# Conclusions

1. First-time simulation of artificial *Fermi*-LAT light curves from magnetic reconnection
2. General characteristics of real LAT light curves are recovered (average flux and minimum variability timescale)
3. To explain the day-long high flux of the brightest flare of 3C 279 is challenging (high external photon density required)
4. Minute-scale bright flares from fast plasmoids are detectable during GTIs
5. Systematic search of minute-scale flares in LAT data on sub-orbital time scales could serve as a test for magnetic reconnection in blazars



If you would like to learn more, take a look at our paper:  
[Meyer, Petropoulou, Christie, 2021, ApJ, 912](#)

# Back-up slides




# Event selection and ROI optimization

- Normalizations left free to vary for sources within 10 deg, all spectral parameters for sources within 5 deg + additional point sources with TS > 25 added iteratively; central source relocalized → yields smooth residual and TS maps
- For weekly and daily light curves: central source spectral parameters and normalizations of sources within 1 deg + isotropic and galactic diffuse normalizations left free to vary.




Data set	Pass 8R3 V6
Event class	P8R3 SOURCE
Energy range, binning	0.1 - 316 GeV, 8 bins per decade
ROI size, binning	15° x 15°, 0.2° per pixel
Zenith angle	< 90°
Time cuts filter	DATA_QUAL>0 && LAT_CONFIG==1; Additionally bright solar flares and GRBs with TS > 100 excised
Fermi tools version / fermipy version	1.23 / 0.19.0
Catalog/s	4FGL, gll_psc_v18.fit
Galactic diffuse template	gll_lem_v07.fit
Isotropic diffuse templates	iso_P8R3_SOURCE_V2_v1.txt

# Theoretical light curves and periodograms for different plasma magnetizations

$\sigma = 3$




$\sigma = 50$



# The observability of plasmoid-powered $\gamma$ -ray flares with the *Fermi* Large Area Telescope

M. Petropoulou\*, M. Meyer, I. Christie on behalf of the Fermi-LAT Collaboration



## What is it about?

Fast  $\gamma$ -ray blazar flares from reconnection and their detectability with Fermi-LAT

## Why is it interesting?



- Origin of fast  $\gamma$ -ray blazar variability is still a mystery!
- Magnetic reconnection models predict fast & luminous  $\gamma$ -ray flares.

## What did we do?

- We created for the first time artificial LAT light curves based on the magnetic reconnection model.
- We searched for detectability of model-predicted features with LAT.

## What did we find?

- General features of real LAT light curves are recovered in the artificial LCs
- Minute-scale flares are detectable with LAT during GTIs



If you would like to learn more, watch our video!